



Fig 1. Chilling Block

Fig 2. In-use Chilling Block



CHILLING BLOCK INSTRUCTIONS FOR USE



SUPPLIES NEEDED

PHARMACY SUPPLIES

- 70% Isopropyl alcohol or equivalent

DESCRIPTION

Chilling Block is a portable chilling device designed to provide cooling of vials and syringes for at least 2 hours for a temperature of -3 °C to 5 °C (27 °F to 41 °F) for the preparation of pharmaceutical products. To attain this temperature, the Chilling Block has to be frozen overnight before each use.

Compartments diameter and depth fit the following components:

- 2 compartments for up-to 100 mL vial
- 1 compartment for up-to 30 mL vial
- 2 compartments for up-to 20 mL syringe

CONTRAINDICATIONS

There are no contraindications for the Chilling Block.

STORAGE AND REUSE

The Chilling Block shall be stored in a cool and dry place.

CLEANING AND DISINFECTION INSTRUCTIONS

1. Clean the Chilling Block after each use. Clean with mild detergent.
2. Disinfect by spraying with 70% Isopropyl alcohol or equivalent and allow to air dry.

WARNINGS AND PRECAUTIONS




Disinfect the Chilling Block using the recommended disinfection instructions and using the recommended cleaning agents.



A frozen Chilling Block may be slippery.



DO NOT USE THE CHILLING BLOCK if the device is damaged or if it leaks.




Do not expose to temperatures below -30 °C (-22 °F) and above 50 °C (122 °F)


INSTRUCTIONS

BEFORE EACH USE:

- 1 Verify that the Chilling Block is not damaged and does not leak.




- 2 12 hours before use place the Chilling Block into a freezer at a temperature of -20 °C to -12 °C (-4 °F to 10.4 °F)



WHEN READY FOR USE:

- 3 Using the handles, take the Chilling Block out of the freezer.



Caution: A frozen Chilling Block may be slippery.



Note: The Chilling Block must be used within 2 hours of removal from the freezer.

EXPLANATION OF SYMBOLS:

REF Catalogue number

SN Serial number

LOT Lot number

Date of manufacture

Caution, consult accompanying documents

Consult instructions for use

Temperature limit

Distributed by:
UroGen Pharma, Inc.
Princeton, NJ 08540

UCB-IFU-001
IFU-Phy-0016592/Ver. 1